

Aureomycin®

Type A Medicated Article

Product Description

Aureomycin® (chlortetracycline) is a Type A Medicated Article, available in granular and meal formulations, for disease control in chickens, including laying hens, turkeys and ducks and for disease treatment in psittacine birds. Aureomycin contains 50, 90 or 100 grams of chlortetracycline activity per pound of premix.

Indications for Use

- For control of infectious synovitis (*Mycoplasma synoviae*) in chickens and turkeys.
- For the control of chronic respiratory disease (CRD) and air sac infection (*Mycoplasma gallisepticum*, *Escherichia coli*) and the reduction of mortality due to *E. coli* in chickens.
- For the reduction in mortality due to *E. coli* infections susceptible to chlortetracycline in chickens.
- For the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) and hexamitiasis (*Hexamita meleagridis*) in turkeys.
- For the reduction of mortality due to paratyphoid (*Salmonella typhimurium*) in turkey poultlets not over 4 weeks of age.
- For the control and treatment of fowl cholera caused by *Pasteurella multocida* in ducks.
- For the treatment of psittacosis caused by *Chlamydia psittaci* in psittacine birds (cockatoos, macaws and parrots).

FDA Status: Category I Drug

- Category I drug and does not require a feed mill license for manufacture of medicated feeds.
- Approved for use in combination with Robenz® (broiler or fryer chickens), Deccox® (chickens), clopidol (chickens), salinomycin sodium (broiler chickens) and monensin (broiler chickens).**

Key Points

- Aureomycin may be used in laying chickens.*
- Broad-spectrum antibiotic.
- Provides protection against a wide range of respiratory and enteric diseases.
- Chilsonation process helps to improve flowability, improve stability and minimize dust.¹
- Part of a broad coccidiosis control and gut health management portfolio, backed by service and coccidiosis expertise to help optimize health and performance.

Storage and Packaging

- Store below 25°C (77°F); excursions permitted to 37°C (99°F).
- Aureomycin is packaged in a 50-pound multiwall paper bag with protective barrier ply.

Withdrawal Information

No withdrawal is required prior to slaughter when used according to label.

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Do not feed to ducks or turkeys producing eggs for human consumption

Poultry Information: Chlortetracycline — Type A Medicated Article

Active Drug Ingredient	Chlortetracycline calcium complex
Ingredient	Dried <i>Streptomyces aureofaciens</i> fermentation product and calcium sulfate

For Use in the Manufacture of Medicated Animal Feeds. For Use in Dry Feed Only. Not for Use in Liquid Medicated Feeds.

Species: CHICKENS

Dose: 100-200 g/ton

Indications for Use: For control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Dose: 200-400 g/ton

Indications for Use: For the control of chronic respiratory disease (CRD) and air sac infection caused by *M. gallisepticum* and *E. Coli* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Dose: 500 g/ton

Indications for Use: For the reduction of mortality due to *E. coli* infections susceptible to chlortetracycline. Feed for 5 days.

Species: PSITTACINE BIRDS

Dose: 10 mg/g of feed

Indications for Use: For the treatment of psittacine birds (parrots, macaws and cockatoos) suspected or known to be infected with psittacosis caused by *Chlamydia psittaci* sensitive to chlortetracycline. Feed continuously for 45 days. Each bird should consume an amount of medicated feed equal to one-fifth of its body weight daily. During treatment, parrots, macaws and cockatoos should be kept individually or in pairs in clean cages.

Caution: Aspergillosis may occur following prolonged treatment.

Warning: Psittacosis, avian chlamydiosis or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals and man. Contact appropriate public health and regulatory officials.

Species: TURKEY POULTS (Not Over 4 Weeks of Age)

Dose: 400 g/ton

Indications for Use: For the reduction of mortality due to paratyphoid caused by *Salmonella typhimurium* susceptible to chlortetracycline.

Species: TURKEYS

Dose: 200 g/ton

Indications for Use: For control of infectious synovitis caused by *M. synoviae* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Dose: 400 g/ton

Indications for Use: For the control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Dose: 25 mg/lb body weight per day

Indications for Use: For the control of complicating bacterial organisms associated with bluecomb (transmissible enteritis; coronaviral enteritis) susceptible to chlortetracycline. Feed continuously for 7 to 14 days.

Species: DUCKS

Dose: 200-400 g/ton

Indications for Use: For the control and treatment of fowl cholera caused by *Pasteurella multocida* susceptible to chlortetracycline. Feed in complete ration to provide from 8 to 28 mg per pound of body weight per day depending on age and severity of disease. Feed for not more than 21 days.

**Aureomycin combined with other drugs may not be approved for use in layers. Refer to 21 CFR 558.128 or consult your Phibro representative.*

Restricted Drug (California) — Use Only as Directed. Not for use in humans. Keep out of reach of children. Approved by FDA under NADA # 048-761
Please refer to the product label for full directions, including mixing directions.

¹ Laing N. Granulated Ingredients: Reducing drug potency. Feed Management. December 1994.

**To learn more about Aureomycin
talk with a Phibro expert at**

800-677-4623 or visit www.pahc.com