

Avatec[®]

Type A Medicated Article

Product Description

Avatec[®] (*lasalocid*) is a Type A Medicated Article for the prevention of coccidiosis in poultry. Avatec contains 90.7 grams (20%) of lasalocid sodium activity per pound of premix.

Indications for Use

- For the prevention of coccidiosis in broiler chickens, growing turkeys and chukar partridges.
- Prevents coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. brunetti*, *E. mivati*, *E. maxima*, *E. meleagritidis*, *E. gallopavonis*, *E. adenoides* and *E. legionensis*.

Key Points

- Avatec provides rotation program advantages for both broiler chickens and turkeys as the only available divalent ionophore.
- Highly effective, broad spectrum and kills at the first stage of development.
- Provides year-round (non-seasonal) coccidiosis protection.
- No depression of feed or water intake.¹

FDA Status

- Category 1 drug and does not require a feedmill license for manufacture of medicated feeds.
- Approved for use in combination with, but not limited to, BMD[®] (broilers, fryers and turkeys) and Lincomix[®] (broilers).*
- If Avatec is used in combination with Lincomix, a VFD is required.

Storage and Packaging

- Store below 25°C (77°F); excursions permitted to 37°C (99°F).
- Avatec is packaged in a 50-pound, multiwall paper bag with protective barrier ply.

Withdrawal Information

- No withdrawal is required prior to slaughter.

See reverse side for complete noncombination use directions, product claims and additional information on this product.

*Reference Title 21 Code of Federal Regulations 558.311 for species-specific combination clearances.

Poultry Information: Lasalocid Sodium — Type A Medicated Article

Active Drug Ingredient	Each pound contains 90.7 g (20%) of lasalocid sodium activity in a carrier suitable for incorporation in feed.		
Mixing Directions	Thoroughly mix the following amounts of Avatec to make one ton of feed to provide from 68 g per ton (0.0075%) to 113 g per ton (0.0125%) lasalocid sodium concentration. When adding less than one pound of premix per ton of feed, it is recommended that the premix first be mixed with a small amount of feed (10 to 25 lb.) prior to incorporation into the final feed.		
For Manufacturing Medicated Poultry and Livestock Feeds Only	Concentration % in Feed	Lasalocid grams/ton	Pounds Avatec® per ton feed
Broiler, Fryer Chickens and Growing Turkeys	0.0075%	68	¾
	0.0100%	90.7	1
	0.01125%	102	1⅛
Broiler, Fryer Chickens, Growing Turkeys, Chukar Partridges	0.0125%	113	1¼
Limitations	Feed continuously as the sole ration.		
Warning	When mixing and handling lasalocid premix, use protective clothing, impervious gloves and a dust mask. Avoid contact with eyes. Operators should wash thoroughly with soap and water after handling.		

Note: In addition to label statements of identity, ingredients and directions for use, labeling of feeds containing Avatec must contain the following statements:

For Broiler or Fryer Chickens Only and/or For Growing Turkeys Only
 For Chukar Partridges (up to 8 weeks of age)
 Only Restricted Drug (California) - Use only as directed.
 Not for human use.
 Feed continuously as the sole ration.

Approved by FDA under NADA # 096-298

¹ Macy LB, et al. Effects of feeding lasalocid on performance of broilers in moderate and hot temperature regimens. *Poult Sci.* 1990;69:1265-1270.

**To learn more about Avatec
 talk with a Phibro expert at**

800-677-4623 or visit www.pahc.com